

**Michigan**  
**Adult Treatment Protocols**  
**NAUSEA & VOMITING**

Date: November 15, 2012

Page 1 of 2

---

## ***Nausea & Vomiting***

### **Pre-Medical Control**

#### **MFR/EMT/SPECIALIST/PARAMEDIC**

1. Follow **General Pre-hospital Care Protocol**.

#### **SPECIALIST/PARAMEDIC**

2. Administer NS IV/IO fluid bolus up to 1 liter, wide open.
3. Hypotensive patients should receive additional IV/IO fluid boluses, as indicated by hemodynamic state. Continue IV/IO fluid bolus to a maximum of 2 liters.

#### **PARAMEDIC**

4. Administer Ondansetron (Zofran) 4mg IV/IM.

### **Post-Medical Control**

5. Repeat Ondansetron (Zofran) 4mg IV/IM.
6. For Pediatric Patients refer to **Pediatric Nausea/Vomiting Protocol**.

MCA Name Delta County  
MCA Board Approval Date 8/8/13  
MDCH Approval Date 9/6/13  
MCA Implementation Date 11/1/13



**Section 1-11**  
*E F Bigsby, MD*

**Michigan  
Adult Treatment Protocols  
NAUSEA & VOMITING**

Date: November 15, 2012

Page 2 of 2

